



**Xelstrym®**  
(dextroamphetamine) **II**  
transdermal system  
4.5mg/9 hours 9mg/9 hours 13.5mg/9 hours 18mg/9 hours

Ready for ADHD treatment

# Your time, Your way

Work with your healthcare provider to determine when to apply and when to remove XELSTRYM.

**XELSTRYM should be removed within 9 hours.**

## What is XELSTRYM?

XELSTRYM is a central nervous system stimulant prescription medicine used for the treatment of Attention-Deficit Hyperactivity Disorder (ADHD) in adults and children 6 years and older. It is not known if XELSTRYM is safe and effective in children younger than 6 years of age.

## IMPORTANT SAFETY INFORMATION

**Abuse, misuse, and addiction:** XELSTRYM is a federally controlled substance (CII) because it contains dextroamphetamine that can be a target for people who abuse prescription medicines or street drugs. Keep XELSTRYM in a safe place to protect it from theft. Never sell or give your XELSTRYM to anyone else because it may cause death or harm to them and it is against the law. Tell your healthcare provider if you or your child have ever abused or been dependent on alcohol, prescription medicines, or street drugs.

**Please read Important Safety Information on pages 14-19 and [click here](#) for Medication Guide and Full Prescribing Information, including the BOXED WARNING.**

Are you ready for

# ADHD treatment Your time, Your way?

**Work with your healthcare provider to determine when to apply and when to remove XELSTRYM®.**

- **XELSTRYM** is the **first and only amphetamine treatment for ADHD** that uses patch technology to deliver medication through the skin\*
- The technology of XELSTRYM allows you and your healthcare provider to tailor your ADHD treatment to meet your daily needs

\*Please see Instructions for Use.

**Please read Important Safety Information on pages 14-19 and [click here](#) for Medication Guide and Full Prescribing Information, including the BOXED WARNING.**

**Xelstrym®**  
(dextroamphetamine) transdermal system  
4.5mg/9 hours 9mg/9 hours 13.5mg/9 hours 18mg/9 hours




Ready for

# Working with your healthcare provider for flexibility with your schedule

**Treatment for ADHD should meet the demands of your changing schedule and offer flexibility.**

With XELSTRYM®, you and your healthcare provider can work together to decide when to apply and when to remove the patch so that your treatment is tailored to your schedule throughout the day.

- Apply XELSTRYM 2 hours before the effect is needed and remove XELSTRYM within 9 hours after application



Are there times of the day when you and your healthcare provider would like to align periods of symptom control with your daily routines?

Please read Important Safety Information on pages 14-19 and [click here](#) for Medication Guide and Full Prescribing Information, including the BOXED WARNING.

**Xelstrym®**  
(dextroamphetamine) <sup>®</sup> transdermal system  
4.5mg/9 hours 9mg/9 hours 13.5mg/9 hours 18mg/9 hours



Meals



Work schedule



Family time



Time with your partner



Social engagements



"Me" time

Ready for

# A treatment you can wear

XELSTRYM® has the same active metabolite (dextroamphetamine) as Vyvanse®\*, a medication that has been used to treat ADHD since 2007.

## How does XELSTRYM work?

XELSTRYM delivers medication that is passively absorbed through the skin to provide continuous drug release into the bloodstream.

## What is transdermal therapy?

Transdermal drug delivery is a method of delivering drugs systemically by applying a drug formulation onto intact and healthy skin.

There are no head-to-head safety or efficacy studies comparing Vyvanse® and XELSTRYM.

\*VYVANSE® is a registered trademark of Takeda Pharmaceuticals U.S.A., Inc.

Please read Important Safety Information on pages 14-19 and [click here](#) for Medication Guide and Full Prescribing Information, including the BOXED WARNING.

**Xelstrym®**  
(dextroamphetamine)   
transdermal system  
4.5mg/9 hours 9mg/9 hours 13.5mg/9 hours 18mg/9 hours



Ready for

# Established safety

## What are the most common side effects of XELSTRYM®?

- |                      |                            |
|----------------------|----------------------------|
| • Decreased appetite | • Increased blood pressure |
| • Headache           | • Muscle twitching (tics)  |
| • Trouble sleeping   | • Vomiting                 |
| • Stomach pain       | • Irritability             |
| • Nausea             | • Increased heart rate     |

Application site reactions have happened while wearing XELSTRYM and after removal of the patch. Symptoms include pain, itching, burning feeling, redness, discomfort, or swelling at the application site. Call your healthcare provider if you or your child develop any application site reactions that do not resolve on their own.


Please read Important Safety Information on pages 14-19 and [click here](#) for Medication Guide and Full Prescribing Information, including the **BOXED WARNING**.

**Xelstrym®**  
(dextroamphetamine)   
transdermal system  
4.5mg/9 hours 9mg/9 hours 13.5mg/9 hours 18mg/9 hours

## A helpful how-to video

### Scan and watch.

Use your mobile device and scan the QR code to watch a video that will show you how to apply your patch.



[Click here to watch the IFU video](#)

Please see Instructions For Use.



Ready for

# Discreet and tailored dosing

**Work with your healthcare provider to tailor your dose to meet your needs.**

Four dosage strengths provide flexible options



Patches, US nickel, and golf ball shown at actual size.

Please read Important Safety Information on pages 14-19 and [click here](#) for Medication Guide and Full Prescribing Information, including the **BOXED WARNING**.

**Xelstrym®**  
(dextroamphetamine)   
transdermal system  
4.5mg/9 hours 9mg/9 hours 13.5mg/9 hours 18mg/9 hours



**10**

## Application wear sites

**Choose from left or right side of the body.**

- Use one patch daily (per 24 hours)
- Do not wear the patch for more than 9 hours
- Change application site when a new patch is applied

Ready for

# A medication you can wear during most of your daily activities

	YES	AVOID
Showering*	✓	
Bathing*	✓	
Swimming*	✓	
Activities that cause sweating†	✓	
Direct heat‡		✗

\*Check to see if the patch has become loose after bathing, showering, or swimming.

†Check the patch fit if you sweat excessively.

‡After applying XELSTRYM®, avoid exposing the application site to direct external heat sources, such as hair dryers, heating pads or electric blankets, heat lamps, saunas, hot tubs, and heated water beds. Exposure to heat can cause too much medicine to pass into your body and cause serious side effects.

Please read Important Safety Information on pages 14-19 and [click here](#) for Medication Guide and Full Prescribing Information, including the BOXED WARNING.

## XELSTRYM Copay Savings Program

Pay as little as \$25 at a Noven C.A.N.™ Partner Pharmacy\*

**Xelstrym®**  
(dextroamphetamine)   
transdermal system  
4.5mg/9 hours 9mg/9 hours 13.5mg/9 hours 18mg/9 hours

Pay As Little As  
**\$25\*** at a Partner Pharmacy

**Xelstrym®**  
(dextroamphetamine)   
transdermal system  
4.5mg/9 hours 9mg/9 hours 13.5mg/9 hours 18mg/9 hours

\*Copay savings offer is available for commercially insured patients. At Noven C.A.N.™ Partner Pharmacies, eligible patients pay as little as \$25 for each XELSTRYM prescription. At non-Partner Pharmacies, eligible patients pay as little as \$50 for each XELSTRYM prescription. Patient is responsible for any remaining balance due after copay savings offer is applied. An offer is available for cash-paying patients.

Care Access Network  
**NOVEN CAN**

Scan the QR code  
to ENROLL TODAY



## What is XELSTRYM?

XELSTRYM is a central nervous system stimulant prescription medicine used for the treatment of Attention-Deficit Hyperactivity Disorder (ADHD) in adults and children 6 years and older. It is not known if XELSTRYM is safe and effective in children younger than 6 years of age.

## IMPORTANT SAFETY INFORMATION

**Abuse, misuse, and addiction: XELSTRYM is a federally controlled substance (CII) because it contains dextroamphetamine that can be a target for people who abuse prescription medicines or street drugs.**

Keep XELSTRYM in a safe place to protect it from theft. Never sell or give your XELSTRYM to anyone else because it may cause death or harm to them and it is against the law. Tell your healthcare provider if you or your child have ever abused or been dependent on alcohol, prescription medicines, or street drugs.

### Do not use XELSTRYM if you or your child are:

- Allergic to amphetamine or any of the ingredients in XELSTRYM. See the end of the Medication Guide for a complete list of ingredients in XELSTRYM.
- Taking or have taken within the past 14 days a medicine called a monoamine oxidase inhibitor (MAOI), including the antibiotic linezolid or the intravenous medicine called methylene blue.

Please read Important Safety Information on pages 14-19 and [click here](#) for Medication Guide and Full Prescribing Information, including the BOXED WARNING.

## IMPORTANT SAFETY INFORMATION (CONTINUED)

### XELSTRYM can cause serious side effects, including:

- **Risks for people with serious heart disease.** Sudden death has happened in people who have heart defects or other serious heart disease. Your healthcare provider (HCP) should check you or your child carefully for heart problems before starting treatment with XELSTRYM. Tell your HCP if you or your child have any heart problems, heart disease, or heart defects. **Call your HCP or go to the nearest hospital emergency room right away if you or your child have any signs of heart problems such as chest pain, shortness of breath, or fainting during treatment with XELSTRYM.**
- **Increased blood pressure and heart rate.** Your HCP should check you or your child's blood pressure and heart rate regularly during treatment with XELSTRYM.
- **Mental (psychiatric) problems.** Including new or worse behavior and thought problems; new or worse bipolar illness; new psychotic symptoms (such as hearing voices, or seeing or believing things that are not real); or new manic symptoms. Tell your HCP about any mental problems you or your child have or about a family history of suicide, bipolar illness, or depression. **Call your HCP right away if you or your child have any new or worsening mental symptoms or problems during treatment with XELSTRYM, especially hearing, seeing or believing things that are not real, or new manic symptoms.**

**Xelstrym®**  
(dextroamphetamine)   
transdermal system  
4.5mg/9 hours 9mg/9 hours 13.5mg/9 hours 18mg/9 hours

## IMPORTANT SAFETY INFORMATION (CONTINUED)

- **Slowing of growth (height or weight) in children.** Children should have their height and weight checked often while on XELSTRYM. Your HCP may stop treatment with XELSTRYM if your child is not growing or gaining height or weight as expected.
- **Circulation problems in fingers and toes (peripheral vasculopathy, including Raynaud's phenomenon).** Fingers or toes may feel numb, cool, or painful. Fingers or toes may change color from pale, to blue, to red. Tell your HCP if you or your child have any numbness, pain, skin color change, or sensitivity to temperature in the fingers or toes. **Call your HCP right away if you or your child have any signs of unexplained wounds appearing on fingers or toes while using XELSTRYM.**
- **Serotonin syndrome.** Symptoms such as agitation, fast heartbeat, flushing, seizures, coma, sweating, loss of coordination, confusion, dizziness, tremors, stiff muscles, muscle twitching, seeing or hearing things that are not real (hallucinations), changes in blood pressure, high body temperature (hyperthermia), nausea, vomiting, or diarrhea may occur. Stop using XELSTRYM and call your HCP or go to the emergency room if symptoms occur. Serotonin syndrome may occur if XELSTRYM is taken with certain medicines and may be life-threatening.
- **Allergic skin rash (contact sensitization).** Stop using XELSTRYM and tell your HCP right away if you or your child develop swelling or blisters at or around the application site.

Please read Important Safety Information on pages 14-19 and [click here](#) for Medication Guide and Full Prescribing Information, including the BOXED WARNING.

## IMPORTANT SAFETY INFORMATION (CONTINUED)

- **Application site reactions.** Reactions have happened while wearing XELSTRYM and after removing it. Symptoms include pain, itching, burning feeling, redness, discomfort, or swelling at the application site. Call your HCP if you or your child develop any application site reactions that do not resolve on their own.
- **New or worsening tics or worsening Tourette's syndrome.** Tell your HCP if you or your child get any new or worsening tics or worsening Tourette's syndrome during treatment with XELSTRYM.

**Before taking XELSTRYM tell your HCP about all the medications you take and any medical conditions including if you:**

- Are pregnant or plan to become pregnant. It is not known if XELSTRYM may harm your unborn baby. There is a pregnancy registry for women who are exposed to XELSTRYM during pregnancy. The purpose of the registry is to collect information about the health of women exposed to XELSTRYM and their babies. For more information on registering with the National Pregnancy Registry for Psychiatric Medications call 1-866-961-2388 or visit online at <https://womensmentalhealth.org/clinical-and-research-programs/pregnancyregistry/othermedications/>.
- Are breastfeeding or plan to breastfeed. XELSTRYM can pass into your milk. Do not breastfeed while taking XELSTRYM. Talk to your doctor about the best way to feed your baby if you take XELSTRYM.

## IMPORTANT SAFETY INFORMATION (CONTINUED)

### What should I avoid while using XELSTRYM?

- Do not drive, operate heavy machinery, or do other potentially dangerous activities until you know how XELSTRYM affects you.
- Avoid exposing the application site to direct external heat sources such as hair dryers, heating pads, electric blankets, heat lamps, saunas, hot tubs, and heated water beds, as exposure to heat can cause too much medicine to pass into your body and cause serious side effects.

### The most common side effects of XELSTRYM include:

- decreased appetite
- trouble sleeping
- stomach pain
- nausea
- increased blood pressure
- headache
- muscle twitching (tics)
- vomiting
- irritability
- increased heart rate

Please [click here](#) to read the Medication Guide and Full Prescribing Information, including the **BOXED WARNING**.

**You are encouraged to report negative side effects of prescription drugs to FDA. Visit <http://www.fda.gov/medwatch> or call 1-800-FDA-1088.**

Please read Important Safety Information on pages 14-19 and [click here](#) for Medication Guide and Full Prescribing Information, including the **BOXED WARNING**.

**Are you ready for  
ADHD treatment**

# **Your time, Your way?**

**Xelstrym<sup>®</sup>**  
(dextroamphetamine) **II**  
transdermal system

4.5mg/9 hours 9mg/9 hours 13.5mg/9 hours 18mg/9 hours

**Work with your healthcare provider to determine  
when to apply and when to remove XELSTRYM.**

**XELSTRYM should be removed within 9 hours.**

For more information, please visit [\*\*xelstrym.com\*\*](https://www.xelstrym.com)

**Please read Important Safety Information on pages 14-19  
and [click here](#) for Medication Guide and Full Prescribing  
Information, including the BOXED WARNING.**

**noven**

XELSTRYM<sup>®</sup> is a trademark of Noven Therapeutics, LLC.

Noven C.A.N. and its related logo are trademarks of Noven Therapeutics, LLC.

©2024 Noven Therapeutics, LLC. All rights reserved. For US audience only.

XEL-3010-16 07/2024